MALACOLOGIA, 2015, 58(1-2): 233-243

THERMAL BIOLOGY OF THE SOUTH AMERICAN APPLE SNAIL ASOLENE PLATAE (CAENOGASTROPODA: AMPULLARIIDAE)

Мапа J. Tiecher, María Е. Seuffert & Pablo В. Martin*

Laboratorio de Ecologia, INBIOSUR (Universidad Nacional del Sur-CONICET)
San Juan 670, 8000, Bahia Blanca, Argentina.

ABSTRACT

Asolene platae is a freshwater snail of the Ampullariidae native from La Plata River basin in South America. Water temperature has an overwhelming influence on the biology of apple snails, though there are no systematic studies related to the thermal biology of this species. The effect oftemperature on the activity, growth and mortality of A. platae was studied through laboratory experiments using constant temperatures of 15, 20, 25, 30 and 35°C. The lower threshold and cumulative degree-days for the embryonic development were estimated under fluctuating temperature. Water temperature had a significant effect on the activity of A. platae, being almost null at 15°C, reaching its maximum in the range 25-30°C and decreasing above 30°C. Regardless of temperature, snails remained resting nearly half of the time, exhibiting less activity than other apple snails across similar temperatures. Mortality was higher for hatchlings than for adults at all temperatures with the exception of 35°C, which resulted in high levels of mortality for both groups. Growth of hatchlings was significantly affected by temperature, being highest between 25 and 35°C. The lower temperature threshold for the embryonic development was around 16°C and the cumulative degree-days to complete devel-opment were estimated between 107.5 and 120.5°C.d. The optimum temperature for A. platae would be 25-30°C. Temperatures of 35°C and above would be harmful, while temperatures below 20°C resulted in rapid decreasing in activity and slower growth rates. The information about the thermal requirements of A. platae will be useful optimizing rearing conditions and developing predictive models of the potential distribution of A. platae in scenarios where it is introduced outside of its native range.
Key words: temperature, growth, mortality, activity, embryonic development; degree-days, threshold.

INTRODUCTION

Ampullariids or apple snails are recognized
worldwide as successful invaders, especially
Pomacea spp. that became serious pests
of aquatic crops (Cowie, 2002; Hayes et al. 2008), promoters of ecosystemic changes in natural wetlands (Carlsson et al., 2004, Fang et al., 2010; Horgan et al., 2014) and vectors of parasitic diseases (Lv et al., 2009). For these reasons, Pomacea has been the target of most research on ampullariids and hence a vast knowledge of its species is available, whereas little is known about the biology and ecology of other apple snail genera like Asolene.
Water temperature has an overwhelming influence on most aspects of the biology of ampullariids, including activity levels, aerial res-piration, growth rate, development, reproduction

*Corresponding author: pmartin@criba.edu.ar

233

and survival (e.g., Freiburg & Hazelwood, 1977; Albrecht et al., 1999; Estebenet & Martin, 2002; Aufderheide et al., 2006; Gettys et al., 2008; Seuffert & Martin, 2009; Seuffert et al., 2010, 2012). However, there are no systematic studies
to date related to the thermal biology of the ge-nus Asolene. The information about the thermal limits of ectotherms in general and aquatic snails in particular would be necessary to understand and predict changes in distribution ranges and abundances in native and invaded areas in the face of global climatic changes (Tomanek, 2008; Madeira et al., 2012; Fordham et al., 2013). Asolene platae (Maton, 1811) is an apple snail native from La Plata River basin, whose range extends from Bolivia to northern Buenos Aires province, Argentina (Cowie & Thiengo, 2003). Recently, this species has been considered a senior synonym of Asolene pulchella (Maton,

234 TIECHER ET AL.

1838) on the basis of their genetic, anatomi-
cal and conchological similarity (Hayes et al.,
2009a; Leon et al., 2014). The embryonic devel-
opment ofA . platae has been determined under
a controlled temperature of 25 + 2°C, extending
for 10.8 days on average (range: 8-16 days; Tiecher et al., 2014). However, the temperature below which embryonic development stops, known as lower temperature threshold (Seuffert et al., 2012), is so far unknown. As temperature increases above the lower threshold, the dura-tion of embryonic development will decrease. However, the combination of temperature above the lower threshold and time required to hatch remains generally constant (Zalom et al., 1983; Seuffert et al., 2012). The estimation of this physiological time (measured in cumula-tive degree-days) and the lower threshold are essential to estimate the duration of embryonic development in thermally fluctuating environ-ments, since a bigger difference between tem-perature and the lower threshold implies an earlier accumulation ofthe required degree-days and hence a faster developmental rate. Among apple snails, these parameters are only known so far for species of Pomacea that have aerial egg masses and calcareous eggshells (Seuffert et al., 2012) but not for genera possessing ge-latinous and subaquatic egg masses such as Asolene (Hayes et al., 2009b).
A series of experiments were designed to study the effect of temperature on the activity of A. platae and the relation with some aspects of its life cycle, such as growth and mortality. Additionally, we estimated the lower threshold and cumulative degree-days for embryonic development. These results provide insights into the fundamental biology of ampullariids and contribute to our understanding of the factors influencing its distribution. These data are also informative for developing predictive models of successful establishment and spread of this species if introduced outside of its native range, and which are critical for assessment of invasion potential by regulatory agencies (e.g., EFSA, 2013, 2014). Furthermore, the results of this study and those on thermal requirements of other ampullariids are useful for optimizing culturing conditions for research and aquacul-ture projects (Seuffert & Martin, 2013).

MATERIALS AND METHODS

Origin and Maintenance of Snails

The snails used in this study were descen-dants of a pool originally obtained from Regatas

pond (Ciudad Autonoma de Buenos Aires, Ar-gentina; 34°33’S, 58°26 W). They were raised in 20 Lcollective aquaria with CaCO; saturated tap water, kept in a room at 25 + 2°C under a 14:10 h (light/dark) photoperiod. The snails were fed with fresh lettuce ad /ibitum and the aquaria were cleaned and renewed weekly. Before each trial, snail shell length (SL, from the apex to the extreme opposite lip of the aperture) was measured with a Vernier caliper to the nearest 0.1 mm.
The laboratory experiments were performed in 10.8 L glass aquaria (20 cm-wide, 27 cm-long and 30 cm-deep). Water temperatures (T) were kept constant with electric thermo-stats located at the bottom of each aquarium to reduce thermal stratification. The aquaria were provided with a 1 cm sand layer to allow the snails to bury themselves. The snails were provided with fresh lettuce as entire leaves and also fragmented in floating particles, allowing pedal surface collecting (Saveanu & Martin, 2013). The aquaria were not provided with artificial aeration.

Experiment 1: Effect of Constant Temperatures on Activity

The water temperatures used in this experi-ment were: 15, 20, 25, 30 and 35°C and for each temperature three snails (SL: 11.1-31.3 mm) were randomly chosen from the general pool. Before the observation period, the snails were acclimated to the target temperature for 48 h. After, they were observed at five minute intervals for over three hours with all treatments run concurrently. At the end of each observa-tion period snails were evaluated to see if they were dead or alive by checking on the heartbeat through the translucent shell or by stimulating them with a probe; in some cases, it was obvi-ous because of the putrefaction state. For each treatment five replicated aquaria were used and each snail was used only once.
During the observation period, the instan-taneous activity or state of each snail was recorded and classified as: feeding (F, consid-ering both shredding and pedal surface collect-ing), crawling (C), ventilating the lung (V, during which it was also registered the number of pumping movements; Seuffert & Martin, 2009), resting (R, on the bottom or clinging to the sides of the aquarium with the foot extended) and inactive (1, with the head and foot fully or partially retracted inside the shell, either lying on the bottom or buried in the sand).
On the basis of the total records during the 3 h observation period, the percentage of time

THERMAL BIOLOGY OF ASOLENE PLATAE „285

spent in the different activities per aquarium was calculated as:

T

number of re
e

cords for ea

ch activity
eee 100
(number of snails per aquarium *
total number of observations)

Experiment 2: Effect of Constant Temperatures on Mortality of Adult Snails and on Mortality and Growth of Hatchlings

The water temperatures used were as in experiment 1. To analyze the effect of tem-perature on the mortality of adult snails, five snails (SL: 14.8-31.5 mm) were randomly as-signed to each temperature. During a 25 day trial period, aquaria were checked every two to three days to determine if snails were alive or dead. Five replicated aquaria were used for each temperature.
To analyze the effect of temperature on the mortality and growth of hatchlings, one week-old hatchlings were reared in tap water in 500 ml plastic bottles containers held inside 10.8 L aquaria containing water maintained at one of the following temperatures: 15, 20, 25, 30 and 35°C. During 20 days the number of dead hatchlings was counted every three days. The five temperatures were replicated three times and the total number of hatchlings per container varied according availability at the start of each trial, but were the same for all five temperatures (31, 21 or 19 for each of the replicates).
At the end of the trials, the mortality (per-centage of dead snails or hatchlings) was calculated for each temperature. A scaled digital photograph was taken of ten randomly chosen surviving hatchlings to determine their shell lengths.

Experiment 3: Threshold Temperature and Degree-days Requirement for Embryonic De-velopment

The egg masses used in this experiment were laid in 2.8 L aquaria by snails reared under the general conditions described previ-ously. Females were allowed to copulate each week to ensure that sperm reserves were not depleted. Due to the lack of external sexual dimorphism, females were recognized while they were laying eggs in a collective aquaria or when an egg mass was found in an individual aquaria where a putative female was isolated after being mounted by a male (Tiecher et al., 2014). Egg masses were incubated in 7 cm

diameter dishes filled with 2 cm of tap water in a room with variable temperature which was recorded hourly with a Hobo® data-logger (T,, °C). Water temperature in the dishes followed closely the air temperature but with a delay of approximately one hour. The egg masses were checked daily to record the appearance of the first hatchling. The counting of hatch-lings continued until the last egg hatched and the remaining eggs were counted to estimate egg mass viability. The duration of embryonic development (d, days) was calculated as the number of days elapsed between the date of egg laying and the date of first hatching. Following the results with Pomacea canali-culata (Lamarck, 1822) reported by Seuffert et al. (2012), the lower temperature threshold (T,) for embryonic development was estimated by three different methods: the least standard deviation from the mean observed number of days (SD-day), the coefficient of variation in days (CV-day) and the linear regression coef-ficient method (REG). The different threshold temperatures were estimated using the equa-tions developed by Yang et al. (1995) based on these methods.
The degree-days (DD, °C.d) for the embryonic development were calculated on the basis of hourly records of temperature by computing the degree-hours and summing over the 24-h period. Degree-hours were calculated as the difference between the recorded hourly mean temperature (T,) and the lower threshold (7,):
DD = > (Th TL)

The cumulative DD were estimated by sum-ming the daily DD over the specific period of development of each egg mass (Seuffert et al., 2012). To estimate the possible error due to the use of air temperatures instead of water tem-peratures we mimicked the smaller variation of the latter using the moving average of the last ten hourly records of air temperature and estimated T, and the cumulative DD again.

Statistical Analysis

The effect of temperature on the snails’ ac-tivity and mortality was investigated through one-way ANOVAs. If homogeneity of variances was rejected by Levene’s test, the dependent variable was arcsine square root transformed and re-analyzed. Means were compared with LSD post hoc test. Differences in hatchlings’ size among treatments were analyzed with nested ANOVAs, with the three replicates (bottles) being the nested random factor and water temperature the main fixed factor.

236 TIECHER ET AL.

100
OResting
О Crawling
Feeding
80] mLung ventilation

O)G Q

(Ts%ip)em net

i>>)

20

FIG. 1. Time spent (%) in different activities by A. platae at different constant tem-peratures (T).

RESULTS

Experiment 1: Effect of Constant Temperatures on Activity

Snails spent nearly half of the total time rest-
ing with the foot extended regardless of tem-perature (range: 41.4-68.8%; Fig. 1), without significant differences among treatments (Table 1). The percentage of time that the snails were inactive (fully or partially retracted) decreased significantly with temperature (Table 1); the mean percentage recorded at 15°C (44%) was significantly higher than the means at 25, 30 and 35°C (Fig. 2), temperatures with very low levels of inactivity (between 0.9 and 7.3%). The time spent ventilating the lung increased signifi-cantly with temperature (Table 1), from zero at 15°C to a maximum of 6.7% at 30°C (Fig. 3). The mean frequency of pumping movements per ventilation event was seven and ranged between 1 and 18. The time spent feeding increased significantly with temperature from 2.5% at 15°C to a maximum of 28% at 25°C, and then decreased significantly from 25 to 35°C (Table 1; Fig. 4). In contrast, the percent-age of time spent crawling was not affected by temperature (Table 1).
At 15 and 20°C the snails were found buried in the sand with a mean frequency of 17.3 and

16.8%, respectively, and at higher tempera-tures (25, 30 and 35°C) burying was almost null (< 1% of total time). During the duration of this experiment, no snails were observed mating and no eggs were laid.

Experiment 2: Effect of Constant Temperatures on Mortality of Adult Snails and on Mortality and Growth of Hatchlings

Temperature had a significant effect on the mortality of adult snails (F4 79 = 4.67, p = 0.022), reaching a maximum of 60% at 35°C (Fig. 5). This mean was significantly higher than the mean mortalities recorded at the other tem-peratures (LSD test, p < 0.05) that were quite

TABLE 1. Summary of one-way ANOVAs for ac-tivity variables of A. platae at different constant temperatures. Dependent variables: time spent inactive (1), ventilating the lung (V), feeding (F), crawling (C) and resting (R); | and V were arcsine square root transformed before the analysis.

| V E C R

Fao 5.335 3.765 8.139 1.410 1.304 р 0.004: > :0:019 0.001 01267 © 01302

THERMAL BIOLOGY OF ASOLENE PLATAE 237.

N 3 100 20 Inactive

> O1 15

10

N O1

T(s%ip)em net

O1©

=)

1520253085
TC)

Ventilating

15:20 2580035
TC)

4
90

40

30

Feeding

15: 20:25: 30.85
тс)

FIGS. 2-4. Means + confidence interval (95%) of time spent in different activities by А. platae at dif-ferent constant temperatures (T); different letters indicate significantly different means after LSD test (y-axis are in different scales). FIG. 2: Inactive; FIG. 3: Ventilating the lung; FIG. 4: Feeding; for the first two variables back-transformed values are shown.

low (6-20%). The mortality of hatchlings was
not significantly affected by temperature (F4 ло
= 3.00, p = 0.072) despite being higher at 35°C
with a mean mortality of 60% as with adults. For
all the other temperatures, mortality was 2.0 to
6.6 times higher for hatchlings than for adults. The mean size of hatchlings after 20 days was significantly affected by temperature (ГР. 135 = 13.774, p < 0.001) while the nested factor (replicated bottles) was not significant (Руд, 135 = 0.895, p = 0.539). The minimum size was recorded at 15°C (mean = 2.42 mm; Fig. 6), be-ing significantly lower than the sizes recorded

80
Q= Hatchlings
—@ Adults

M(o%r)t aN&aOlO)=d=a ia)l lt y

al

15 20 25 30 35
TEC)

FIG. 5. Means + confidence interval (95%) of the mortality (percentage of dead snails) of adults and hatchlings of A. platae at different constant tem-peratures (T); different letters indicate significantly different means for adults after LSD test.

at all the other temperatures (LSD test, p < 0.05). The biggest size was obtained at 30°C, being 48.7% higher than the minimum and significantly bigger than the size recorded at 20°C but not significantly different from those of 25° or. 35°C.

Experiment 3: Threshold Temperature and Degree-days Requirement for Embryonic De-velopment

A total of 84 egg masses were used in this experiment. The mean viability of egg masses

FIG. 6. Means + confidence interval (95%) of shell length (SL) of hatchlings of A. platae attained after 20 days at different constant temperatures (T); different letters indicate significantly different means after LSD test.

238 TIECHER ET AL.

— — А platae (a)
mms 2 canaliculata
=== A. platae (b)
«== M. cornuarietis (a)
: M. cornuarietis (b)
«fa | anistes ovum

Dd(due)rv aetlionp ment

e el bol er ee iggy. Bo) ran 12 67 1A@E0S GO 1139 TC)

FIG. 7. Duration of embryonic development of A. platae at different temperatures (T) [a: calculated with the extreme estimations obtained in this paper (Table 2); b: reported by Tiecher et al. (2014)], durations of two other species with subaquatic egg masses [Marisa cornuarietis (a: Demian & Yousif, 1973, b: Aufderheide et al., 2006) and Lanistes ovum (Heiler et al., 2007)] and those obtained for aerial egg masses of Pomacea canaliculata with the same methods used here (bars indicates ranges of temperatures and/or durations).

(100 * number of live hatchlings / total number
of eggs) was 57.8% and fluctuated between
1.1 and 100%. For the estimations of thresh-
olds and degree-days we decided to use the
data obtained from egg masses with viabilities greater than 10%, leaving a total of 77 egg masses. The duration of embryonic develop-ment (d) varied from a minimum of 9 days to a maximum of 26 days (mean + SD = 14.57 + 3.18 d). The estimations of the lower tem-

TABLE 2. Threshold temperatures (°C) and cumu-lative degree-days (DD; °C.d) for the embryonic development of A. platae. Threshold temperatures were estimated by different methods: the least standard deviation in days (SD-day), the coef-ficient of variation in days (CV-day) and the linear regression coefficient method (REG); the values for the cumulative DD are mean + SD.

Method Threshold Cumulative DD

SD-days 16.1 114422354
CV- days 13:7 120.7 + 24.5
REG 16.6 107 6 +219

perature threshold for the embryonic develop-ment of A. platae varied between 15.7°C and 16.6°C according to the method used (Table 2), and the corresponding mean cumulative DD estimations ranged between 107.6°C.d and 120.7°C.d. The cumulative DD of each egg mass was not related to the mean temperature to which it was exposed during development (F у тв = 0.029, р = 0.864, estimated slope + SD = -0.51 + 2.97) indicating that degree days were constant within the temperature range to which the eggs were exposed (19-28°C). The estimations obtained using the moving average of water temperatures (whose greater delay reduced the variation in 4.5%) were only 1.5% higher and 2.4% lesser at most for the lower threshold and cumulative degree-day estima-tions, respectively.

DISCUSSION

Water temperature had a significant effect on the general level of activity of Asolene platae. Regardless of temperature, snails remained resting attached to the substratum nearly half

THERMAL BIOLOGY OF ASOLENE PLATAE 239

of the time. At 15°C snails were inactive (i.e., fully or partially retracted) the other 44% oftime; only during 1% of the time snails were found crawling orfeeding. The general level of activity ofA . platae reached its maximum in the range 25-30°C, and the time that the snails spent buried was almost null above 25°C. Above 30°C activity began to decrease, increasing the percentage of time resting.
The level of inactivity at 15°C comprised near-ly half of the total observation time. In a study of the thermal biology of P canaliculata, it was reported that although the time inactive was lower at 15°C (25%) it steeply increased to 98% at 10°C (Seuffert et al., 2010). These results suggest that A. platae will also be inactive at 10°C. On the other hand, A. platae snails spent nearly half of the time resting, with the foot extended but motionless, regardless of water temperature. Based on the scarce information available, A. plate is apparently a slow growing (MJT, unpublished results) species, much less mobile than P. canaliculata which spends only 26% of the time on average resting between 15 and 30°C (Seuffert et al., 2010).
The time spent feeding increased significantly with temperature, reaching its maximum at 25-30°C and decreasing at 35°C. This increase in feeding corresponded with the decrease of inactivity above 20°C and not to a decrease in resting time as it was reported for Р canaliculata (Seuffert et al., 2010). Feeding in this species was maximum at 25°C and began to decrease with higher temperatures. The level of feeding of Pomacea maculata Perry, 1810, was mini-mum at 15°C and reached its maximum at 30°C (Gettys et al., 2008). For ectothermic animals, this increase seems to be essential to fulfill the energetic demands of an increased metabolism. The decrease in feeding recorded at 35°C may indicate that temperatures above 30°C begin to be suboptimal and even harmful for A. platae in the midterm, since growth decreases and mortality reaches its maximum at 35°C.
The amount of time that A. platae individu-als spent crawling (average 17.5%) was not affected by temperature. A similar result was reported for P. canaliculata, which spent almost the same proportion of time crawling (20.2%) regardless of temperature variation (Seuffert et al., 2010). As it was previously proposed, a plausible explanation is that the snails can crawl faster with increasing temperature. In consequence, they do not spend additional time to move between different places to perform their routine activities.

Ampullariids have both a functional gill and a lung and are considered as amphibious snails (Andrews, 1965; Berthold, 1991; Seuffert 8 Martín, 2009). The maximum level of lung ventilation in A. platae was recorded at 30*C, probably related to an increase in metabolic rate (Akerlund, 1969; Freiburg & Hazelwood, 1977) and to the fact that the levels of dissolved oxygen decrease with increasing temperatures, since the aquaria were not artificially aer-ated. With increasing levels of hypoxia, the significance of aerial respiration increases, becoming a complementary mechanism to branchial intake of oxygen (Seuffert & Martín, 2010). Whereas Р canaliculata breathes air between 15 and 35°C, irrespective of water aeration (Seuffert & Martin, 2009), A. platae ventilates the lung only between 25 and 35°C in non-aerated water. This indicates a lower degree of dependence on aerial respiration in A. platae, which is correlated with the smaller size of its siphon (Seuffert & Martin, 2009) and the late development of its lung (Tiecher et al., 2014). The number of pumping movements per ventilation event varied greatly among rep-licates and showed no relation to temperature. Similarly, no effect oftemperature was detected on the number of pumping movements of Р canaliculata (Seuffert & Martin, 2009). Mortality was higher for hatchlings than for adults at all temperatures with the exception of 35°C, which resulted in high levels of mortality for both groups. Temperature significantly af-fected the mortality of A. platae adults but no differences were detected with the mortality of hatchlings. By contrast, an effect of tempera-ture on the mortality rate of hatchlings was re-ported for Marisa cornuarietis (Linnaeus, 1758) and P canaliculata (Selck et al., 2006; Seuffert & Martin, 2013). The influence of temperature on the mortality rates of A. platae hatchlings was probably masked by their high intrinsic mortality rates (MJT, unpublished data), since the survivorship curves of A. platae generally conform to Type Ш (Deevey, 1947). A different pattern has been reported for P. canaliculata (survivorship curves Type |; Estebenet & Mar-tin, 2002), which evidences a very low intrinsic mortality of hatchlings and juvenile snails (Es-tebenet & Cazzaniga, 1992, 1998), even when exposed to extreme temperatures (only 20% after ten weeks of exposure at 35°C; Seuffert & Martin, 2013).
The shell length of hatchlings attained after 20 days was significantly affected by tempera-ture, being highest at 25, 30 and 35°C. The

240 TIECHER ET AL.

smaller snails were obtained in the treatment
of 15°C. Similarly, the maximum shell length
of Р canaliculata hatchlings after ten weeks
was recorded between 25 and 35°C, being significantly higher than the sizes attained
with 15 and 20°C (Seuffert & Martin, 2013). Hatchlings of P. maculata reared at 15°C did not grow at all and at 35°C grew significantly less than at 20-30%C (Gettys et al., 2008). The growth rate of M. cornuarietis (Linnaeus, 1758) juveniles significantly decreased with temperature from 28°C to 22°C (Aufderheide et al., 2006; Selck et al., 2006). Based on this information we conclude that the optimum tem-perature range for growth varies a little among these South American ampullariids, but for all of them temperatures below 20°C will result in a stunted growth.
The risks of predation, severe hypoxia or prolonged air exposure of subaquatic egg masses (McMahon, 1983; Turner et al., 2007; Tiecher et al., 2014) increase with the dura-tion of development for most oviparous snails inhabiting freshwater habitats, which are highly variable and unpredictable as compared with marine ones. Temperature and oxygen are the main environmental drivers of the duration of development although the effects of tempera-ture may be contradictory, increasing mitotic rates but also reducing oxygen concentrations in water. The analyses of this trait would be important to understand the selective pressures that aquatic snails faced during colonization of inland waters from the sea or terrestrial habi-tats (McMahon, 1983; Dillon, 2000) and also the evolution of aerial egg masses in some lineages (Hayes et al., 2009b).
The duration of embryonic development of A. platae predicted by our degree-day mod-els at different temperatures was compared to the durations previously reported for this species and for other species of ampullariids with subaquatic egg masses and also to the duration estimated for the aerial egg masses of P. canaliculata using the same methods (Fig. 7). Due to the small size and flat shape of the dishes used for the incubation of egg masses, the effect of using air temperature instead of water temperature in our present estimations was smaller than the effect of the three different calculation methods. The durations reported by Tiecher et al. (2014) for egg masses incubated under controlled temperature (25 + 2°C) in the same laboratory and deposited by snails from the same pool as the present study showed a wide variation but the mean duration (10.8

days) agrees with the predictions from our degree-day models. The lower temperature threshold for the embryonic development of A. platae is around 16*C, coinciding with the values of 15.7-16.2°C estimated for P. canaliculata. The cumulative degree-days estimated for A. platae ranged between 107.5 and 120.5°C.d, being also very close to those obtained for P canaliculata (120.7-132.2°C.d.; Seuffert et al., 2012; Fig. 7). On the other hand, up to 1400°C.d have been estimated for acroloxid limpets inhabiting a high latitude lake (Shirokaya & Röpstorf, 2003).
The duration of development in the tem-perature range of 25-30°C for M. cornuarietis (Demian & Yousif, 1973), whose subaquatic egg masses are similar to those of A. platae (Tiecher et al., 2014), are quite coincident with our estimations. In contrast, the duration of 20 days reported for M. cornuarietis in the range of 15-20°C is much shorter than that predicted by our model as temperature ap-proaches the lower threshold (30 days below 20°C). Even though it is a different species, it seems odd that development occurs at such low temperatures, given the tropical origin of the snails used in that study (Puerto Rico). Demian & Yousif (1973) only mentioned the extreme temperatures recorded (15-20°C) so the actual temperature regime under which the eggs developed is unknown. In another study, M. cornuarietis eggs took eight days at 28°C and 17 days at 22°C to hatch (Aufderheide et al., 2006), which is consistent with the reports for this species in the range 25-30°C and only two to three days shorter than our estimates for A. platae. In contrast, the only report to date of the development of an Old World ampullariid with subaquatic egg masses (Lanistes ovum Peters, 1845) indicates that it requires 6-7 days at 26-28°C (Heiler et al., 2007), which is faster than the development for А. platae and M. cor-nuarietis at similar temperatures. Perhaps this is related to the fact that the eggs of L. ovum are disposed in a single layer and hence there is a reduced chance of competition for oxygen within the gelatinous egg mass, which usually delays hatching (Moran & Woods, 2007; Lar-dies & Fernändez, 2002). On the other hand, the cleidoic eggs of P. canaliculata develop at all temperatures at a slightly lower rate (Seuffert et al., 2012) than the aquatic eggs of A. platae studied here. However, the difference may not be significant when taking into account the high inter- and intra-egg mass variation in the duration of development in A. platae (Tiecher

THERMAL BIOLOGY OF ASOLENE PLATAE 241

et al., 2014) and the consequently high stan-dard deviations in the degree-days estimations (Table 2).
Considering the overall results found in this study, we conclude that the optimum tem-perature for A. platae populations would be the range 25-30°C. These temperatures were the optimum for the performing of this species’ routine activities. Temperatures of 35°C and above would be harmful for the snails since not only the levels of activity dropped above 30°C but also mortality rates were higher. On the other hand, at temperatures of 20°C and below the level of general activity rapidly de-creased and the snails grew at a slower rate. Asolene platae exhibits a wide range of thermal tolerance, similar to that of P canaliculata from Encadenadas del Oeste basin (Seuffert et al., 2010). However, the lower activity levels at low temperatures and the higher mortalities at high temperatures probably explains the narrower latitudinal range on A. platae (25 to 35°S; Cas-tellanos & Fernandez, 1976) as compared to P canaliculata (20 to 38°S; Hayes et al., 2012). Even though A. platae is not regarded as an invasive species, it is entering in the aquarium pet trade and thus there are chances of dis-persal to other regions of America and other continents by this mean in the future (Smith, 2006). Specimens are being obtained from natural populations within the natural range of A. platae (Lower Parana river and the southern shore of La Plata river) and offered as Zebra Apple Snails in pet shops and in aquarist web pages (PRM; personal observations). Temperature has been considered the main determinant of establishment in models predict-ing the spread of Pomacea spp. (e.g., Lv et al., 2009; Byers et al., 2013; EFSA, 2013). Based only on the thermal regime, the potential range of distribution of A. platae in the possible sce-nario of an introduction would be less extensive than the currently invaded regions with Poma-cea spp. (southern and eastern Asia, southern USA, Pacific Islands and Spain; Rawlings et al., 2007; Hayes et al., 2008; Lopez et al., 2010), nevertheless it could include many temper-ate and subtropical regions around the world. However, other factors like predation, competi-tion and feeding need to be considered when developing predictive models of the success of its establishment outside of its native range. Knowing the potential range of distribution of A. platae would be useful for detecting and control-ling incipient invasions of these apple snails.

ACKNOWLEDGMENTS

This work was funded with grants by CONICET (“Consejo Nacional de Investigaciones Cientifi-cas y Técnicas”, PIP 112 200901 00473), UNS (“Universidad Nacional del Sur’, PGI 24/B185) and ANPCyT (“Agencia Nacional de Promociön Cientifica y Tecnolögica, PICT 2012-1956). MES and PRM are researchers in CONICET.

LITERATURE CITED

ÄKERLUND, G., 1969, Oxygen consumption of the ampullariid snail Marisa cornuarietis L. in relation to body weight and temperature. Oikos, ааа, A
ALBRECHT, Е. A., М. В. CARRENO & A. CAS-TRO-VAZQUEZ, 1999, A quantitative study of environmental factors influencing the seasonal onset of reproductive behaviour in the South American apple-snail Pomacea canaliculata (Gastropoda: Ampullariidae). Journal of Mol-luscan Studies, 65: 241-250.
ANDREWS, E. B., 1965, The functional anatomy of the mantle cavity, kidney and blood system of some pilids gastropods (Prosobranchia). Journal of Zoology, 146: 70-94. AUFDERHEIDE, J., R. WARBRITTON, N. POUNDS, S. FILE-EMPERADOR, C. STAPLES, N. CASPERS & V. FORBES, 2006, Effects of husbandry parameters on the life-history traits of the apple snail, Marisa cornuarietis: effects of temperature, photoperiod, and population density. Invertebrate Biology, 125(1): 9-20. BERTHOLD, T., 1991, Vergleichende Anatomie, Phylogenie und Historische Biogeographie der Ampullariidae (Mollusca: Gastropoda). Abhan-dlungen des Naturwissenschaftlichen Vereins in Hamburg Series, (NF), 29: 1-256.
BYERS. JEW. ©. MCDOWELL, Э.А. DODD, R. S. HAYNIE, L. M. PINTOR & S. B. WILDE, 2013, Climate and pH predict the potential range ofthe invasive apple snail (Pomacea insularum) in the southeastern United States. PLoS One, 8: 886812. :
CARLSSON, N. O., C. BRONMARK & L. A. HANSON, 2004, Invading herbivory: the golden apple snail alters ecosystem functioning in Asian wetlands. Ecology, 85(6): 1575-1580. CASTELLANOS, Z. J. À. & D. FERNANDEZ, 1976, Gastropoda Ampullariidae. Pp. 1-33, in: К. А. RINGUELET, ed., Fauna de agua dulce de la Repüblica Argentina, FECIC, Buenos Aires. COWIE, R. H., 2002, Apple snails (Ampullariidae) as agricultural pests: their biology, impacts and management. Pp. 145-192, in: G. M. Barker, ed., Molluscs as crop pests. CABI Publishing, Wallingford, New Zealand, 468 pp.
COWIE, К. Н. & S.C. THIENGO, 2003, The apple snails of the Americas (Mollusca: Gastropoda: Ampullariidae: Asolene, Felipponea, Marisa,

242 TIECHER ET AL.

Pomacea, Pomella): a nomenclatural and type
catalog. Malacologia, 45(1): 41-100.
DEEVEY, E. S., 1947, Life tables for natural popu-
lations of animals. Quarterly Review of Biology,
22: 283-314.
DEMIAN, E. & Е. YOUSIF, 1973, Embryonic devel-opment and organogenesis in the snail Marisa cornuarietis (Mesogastropoda: Ampullariidae). Malacologia, 12(1): 123-150.
DILLON, R. T., 2000, The ecology of freshwater molluscs. Cambridge University Press, 509 pp.
EFSA [European Food Safety Authority, Panel on Plant Health], 2013, Scientific Opinion on the assessment of the potential establishment of the apple snail in the EU. EFSA Journal 2013, 11(12): 3487, 49 pp.
EFSA [European Food Safety Authority, Panel on Plant Health], 2014, Scientific Opinion on the environmental risk assessment ofthe apple snail for the EU. EFSA Journal 2014, 12(4): 3641, 97 pp.
ESTEBENET, A. L. & N. J. CAZZANIGA, 1992, Growth and demography of Pomacea canal-iculata (Gastropoda: Ampullariidae) under laboratory conditions. Malacological Review, РЕН
ЕЗТЕВЕМЕТ, A. L. 8 М. J. CAZZANIGA, 1998, Sex related differential growth in Pomacea canaliculata (Gastropoda: Ampullariidae). Jour-nal of Molluscan Studies, 64: 119-123. ESTEBENET, A. L. 4 P. R. MARTIN, 2002, Poma-cea canaliculata (Gastropoda: Ampullariidae): Life-history traits and their plasticity. Biocell, 26: 83-89.
FANG, L., P. K. WONG, L. LIN, C. LAN 4 J. W. QIU, 2010, Impact of invasive apple snails in Hong Kong on wetland macrophytes, nutrients, phytoplankton and filamentous algae. Freshwa-ter Biology, 55: 1191-1204.
FORDHAM, D. A., С. MELLIN, В. D. RUSSELL, К. H. AKCAKAYA, С. J. A. BRADSHAW, М. E. AIELLO-LAMMENS, J. М. CALEY, $. D. CON-NELL, S. MAYFIELD, S. A. SHEPHERD & B. W. BROOK, 2013, Population dynamics can be more important than physiological limits for determining range shifts under climate change. Global Change Biology, 19 (10): 3224-3237. FREIBURG, M. W. & D. H. HAZELWOOD, 1977, Oxygen consumption of two amphibious snails: Pomacea paludosa and Marisa cornuarietis (Prosobranchia: Ampullariidae). Malacologia, 16: 541-548.
GETTYS, L. A., W. T. HALLER, С. В. MUDGE $ T. J. KOSCHNICK, 2008, Effect of temperature and feeding preference on submerged plants by the island apple snail, Pomacea insularum (d’Orbigny, 1839) (Ampullariidae). The Veliger, 50: 248-254.
HAYES, К. A., К. H. COWIE, A. JORGENSEN, Ri СНЕ. С. ALBRECHT & S.C. THIENGO, 2009b, Molluscan models in evo-lutionary biology: apple snails (Gastropoda: Ampullariidae) as a system for addressing fundamental questions. American Malacological Bulletin, 27: 47-58.

HAYES, К. А., К. H. COWIE 8 $. С. THIENGO, 2009a. А global phylogeny of apple snails: Gondwanan origin, generic relationships, and the influence of outgroup choice (Caenogas-tropoda: Ampullariidae) Biological Journal of the Linnean Society, 98, 61-76.
HAYES, К. A., К. H. COWIE, $. С. THIENGO & E. E. STRONG, 2012, Comparing apples with apples: clarifying the identities of two highly invasive Neotropical Ampullariidae (Caeno-gastropoda). Zoological Journal of the Linnean Society, 166: 723-753.
HAYES, K.A., К. С. JOSHI, $. С. THIENGO & К. Н. COWIE, 2008, Out of South America: multiple origins of non-native apple snails in Asia. Diver-sity and Distributions, 14: 701-712.
HEILER, К. С. М., В. SCHULTHEIR, T. GEERTZ 8 С. ALBRECHT, 2007, Caught in the act: insights into the sex life of Lanistes ovum (Gastopoda: Ampullariidae). Antwerp: World Congress of Malacology, Abstracts, p. 94.
HORGAN, F. G., A. M. STUART & E. P. KUDA-VIDANAGE, 2014, Impact of invasive apple snails on the functioning and services of natural and managed wetlands. Acta Oecologica, 54: 90-100.
LARDIES, M.A. & M. FERNANDEZ, 2002, Effect of oxygen availability in determining clutch size in Acanthia monodon. Marine Ecology Progress Series, 239: 139-146.
LEON, J.A., К.А. HAYES & R. H. COWIE, 2014, Ampullariidae revisionary systematics: claryify-ing the taxanomic status of Asolene platae (Ma-ton, 1811) and Asolene pulchella (Anton, 1838). Mollusca 2014. The meeting of the Americas (Abstracts), Mexico City, Mexico, 119-120. LOPEZ, M. A., С. В. ALTABA, К. В. ANDREE € V. LOPEZ, 2010, First invasive apple snails in Europe. Tentacle, 18: 26-28.
LV, S., Y. ZHANG, H. X. LIU, L. HU, К. YANG, P. STEINMANN, Z. CHEN, L. Y. WANG, J. UTZ-INGER & X. N. ZHOU, 2009, Invasive Snails and an Emerging Infectious Disease: Results from the First National Survey on Angiostrongylus cantonensis in China. PLoS Neglected Tropi-cal Disease 3(2): e368. doi:10.1371/journal. pntd.0000368.
MADEIRA, D., L. NARCISO, Н. N. CABRAL & C. VINAGRE, 2012, Thermal tolerance and poten-tial impacts of climate change on coastal and estuarine organisms. Journal of Sea Research,
70. 92-41.
MATON, W. G., 1811, Description of seven new species of Testacea. Transactions of the Lin-nean Society of London, 10: 325-332, pl. 2. McMAHON, БК. F., 1983, Physiological ecology of freshwater pulmonates. Pp. 359-430, in: W. D. RUSSELL-HUNTER, ed., The Mollusca VI, Ecology. Academic Press, New York, 697 pp.
МОКАМ, А. L. & H. A. WOODS, 2007, Oxygen in egg masses: interactive effects of temperature, age, and egg-mass morphology on oxygen sup-ply to embryos. The Journal of Experimental ВОИ 220 172271:
RAWLINGS, T. A., K. A. HAYES, R. H. COWIE & T. M. COLLINS, 2007, The identity, distribution,

THERMAL BIOLOGY OF ASOLENE PLATAE 243

and impacts of non-native apple snails in the continental United States. BMC Evolutionary Biology, 7: 97.
SAVEANU, L. & P. R. MARTIN, 2013, Pedal surface collecting as an alternative feed-ing mechanism of the invasive apple snail Pomacea canaliculata (Caenogastropoda: Ampullariidae). Journal of Molluscan Studies, To: 11-122
SELCK, Н., у. AUFDERHEIDE, NePOUNDS, С. STAPLES, N. CASPERS & V. FORBES, 2006, Effects of food type, feeding frequency, and temperature on juvenile survival and growth of Marisa cornuarietis (Mollusca: Gastropoda). Invertebrate Biology, 125(2): 106-116. SEUFFERT, M. E. & P. В. MARTIN, 2009, Influ-ence of temperature, size and sex on aerial respiration of on Pomacea canaliculata (Gas-tropoda: Ampullariidae) from southern Pampas, Argentina. Malacologia, 51: 191-200. SEUFFERT, M. E. & P. R. MARTIN, 2010, De-pendence on aerial respiration and its influence on microdistribution in the invasive freshwater snail Pomacea canaliculata (Caenogastropoda, Ampullariidae). Biological Invasions, 12: 1695-1708. р
SEUFFERT, М.Е. &Р. В. MARTIN, 2013, Juvenile growth and survival ofthe apple snail Pomacea canaliculata (Caenogastropoda: Ampullariidae) reared at different constant temperatures. SpringerPlus, 2: 312.
SEUFFERT, М. E., $. BUREL8A P. В. MARTÍN, 2010, Influence ‘of water temperature on the activity of the freshwater snail Pomacea canali-culata (Caenogastropoda: Ampullariidae) at its southernmost limit (Southern Pampas, Argen-tina). Jounal of Thermal Biology, 35: 77-84. SEUFFERT, М. E., L. SAVEANU & P. В. MARTÍN, 2012. Threshold temperatures and degree- day estimates for embryonic development of the invasive apple snail Pomacea canaliculata

(Caenogastropoda: Ampullariidae). Malacolo-gia, 55(2): 209-217.
SHIROKAYA, А. А. & P. RÖPSTORF, 2003, Morphology of syncapsules and the duration of embryogeny of Baikalian endemic limpets (Gastropoda, Pulmonata, Acroloxidae). Berliner Paläobiologische Abhandlungen, 4: 183-192. SMITH, J. W., 2006, Ampullariidae pathways. Pp. 113-120, in: К. С. JosHi & L. S. SEBASTIAN, eds., Global advances in ecology and management of golden apple snails. Philippine Rice Research Institute, Nueva Ecija, Philippines, 600 pp. TIECHER, M. J., S. BURELA & P. R. MARTIN, 2014, Mating behavior, egg laying, and em-bryonic development in the South American apple snail Asolene pulchella (Ampullariidae, Caenogastropoda). /nvertebrate Reproduction & Development, 58(1): 13-22.
TOMANEK, L., 2008, The importance of physi-ological limits in determining biogeographical range shifts due to global climate change: the heat shock response. Physiological and Bio-chemical Zoology, 81(6): 709-717.
TURNER, A. M., R. R. TURNER & S. R. RAY, 2007, Competition and intraguild egg preda-tion among freshwater snails: re-examining the mechanism of interspecific interactions Oikos, 116 (11) 1895=1903:
YANG, S., J. LOGAN & D. L. COFFEY, 1995, Mathematical formulae for calculating the base temperature for growing degree days. Agricul-tural and Forest Meteorology, 74: 61-74. ZALOM, F. G., P. B. GOODELL, L. T. WILSON, W. W. BARNETT & W. J. BENTLEY, 1983, De-greedays: the calculation and use of heat units in pest management. Leaflet 21373. Division of Agriculture and Natural Resources, University of California, Davis, California, U.S.A., 10 pp.

Revised ms. accepted November 5, 2014